

Topical review

Fear-avoidance model of chronic musculoskeletal pain: 12 years on

Johan W.S. Vlaeyen^{a,b,*}, Steven J. Linton^c^a Research Group on Health Psychology, Department of Psychology, University of Leuven, Leuven 3000, Belgium^b Department of Clinical Psychological Science, Maastricht University, Maastricht, The Netherlands^c Center for Health and Medical Psychology, Örebro University, Örebro, Sweden

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

1. Introduction

It is widely acknowledged that pain is a universal experience that affects human beings across the life span, serving an important protective function. Typical protective behaviors are the withdrawal from the noxious stimulus, nonverbal expressions that signal others for impending harm, and verbal utterances. Some of these occur involuntarily, as a reflex, whereas other behaviors are more deliberate. However, there is accumulating evidence that it is not pain itself, but the meaning of pain that predicts the extent to which individuals engage in these protective behaviors [1,3]. About a decade ago, we summarized the research evidence supporting the role of fear of pain in the development of chronic pain disability, presented a model incorporating basic mechanisms, but also noted a number of unresolved issues that called for further scientific attention [39] (Fig. 1). In the last decade, the number of studies on this subject has increased exponentially [21], and novel directions are being proposed [6]. Two main stances have emerged. First, although pain has intrinsic threatening features, the threat value of similar pain stimulus may vary across contexts and individuals. Second, protective responding may be adaptive in the short term, but may paradoxically worsen the problem in the long term. In the current updated review, we briefly summarize the progress made since, and highlight a selected number of remaining challenges and areas for future research.

2. The pain-related fear–disability association

In our 2000 review, the fear-avoidance (FA) model was introduced as a way to describe how pain disability, affective distress, and physical disuse develop as a result of persistent avoidance behaviors motivated by fear. The FA model has become increasingly popular, and a large body of evidence is in line with its

assumptions. Research supporting the FA model stems from cross-sectional studies with chronic pain patients [21], prospective studies in acute pain [10,17,33] and studies using structural equation modeling examining the dynamic and sequential relationships among the variables of the FA model [11,40]. Collectively, these findings underscore the important role of pain-related fear in the development of disability. One of the unanswered questions, however, is how pain-related fear occurs in the first place.

3. The acquisition of pain-related fear

By virtue of its biological significance, pain is an important motivator in learning. Indeed, pain informs the individual that there is the imminent or actual threat of body damage. Therefore, pain is considered an unconditioned stimulus (US) that activates an immediate defensive system. Fostering successful adaptation, individuals subsequently anticipate the occurrence of a US by gathering propositional knowledge about the association between neutral cues or conditioned stimuli (CS) and the US. In Pavlovian conditioning, a conditioned response (CR) is elicited when the individual is exposed to a CS, also in the absence of the US. What kind of stimuli are involved in such learning, and what are the potential sources of information leading to such propositional knowledge (Fig. 2)?

Because of their relative ease to use, most fear learning researchers employ exteroceptive (mostly visual) stimuli. For example, when one color (CS+) signals the presentation of painful stimulus (US), and another color does not (CS–), participants usually respond with greater potentiated eye-blink startle, heightened skin conductance, and cardiac deceleration in the presence of the CS+, as compared to CS–, in the absence of the US [4]. In clinical situations, however, and by virtue of their functional proximity, interoceptive and proprioceptive stimuli, rather than exteroceptive ones, may be better predictors of pain as a US. Interoceptive stimuli are those that provide afferent information from receptors that monitor the internal state of the body [8]. Interoceptive fear conditioning therefore occurs when an association between and interoceptive CS and a US has been established [7]. Proprioception is restrictively defined as the perception of posture and movement, also referred to as postural somesthesia [20]. Proprioceptive fear conditioning is particularly relevant in patients with pain in the musculoskeletal system. A recent study used joystick movements, of which the direction predicted painful

* Corresponding author at: University of Leuven, Research Group on Health Psychology, Department of Psychology, Leuven 3000, Belgium. Tel.: +3 216325915; fax: +3 216325923.

E-mail address: johan.vlaeyen@psy.kuleuven.be (J.W.S. Vlaeyen).




Fig. 1. Graphical display of the fear-avoidance model, reproduced from Vlaeyen and Linton [39].




Fig. 2. Graphical display of the acquisition and extinction of pain-related fear, including stimulus topography, protective behaviors and possible pathways. US: Unconditioned stimulus, CS: Conditioned stimulus, CR: Conditioned response.

shock to the hand (e.g., moving upward as CS+ and moving downward as CS-) [26]. As compared with a condition in which both movements were explicitly unpaired with painful shock, the CS+ movement elicited increased fear of movement-related pain, larger eye-blink startle amplitudes, and slower movement latency responses than the CS-, validating the acquisition of fear of movement-related pain in healthy individuals.

Regardless of stimulus topography, at least 3 distinct pathways for acquiring knowledge between potential cues and pain have been proposed. First, people can learn from direct experiences, as was the case in the abovementioned studies [4,26]. However, there are also indirect pathways, such as transmission through verbal instructions or observation. Verbally transmitted information can hold semantically negative information that may yield relevant information about the relationship between 2 stimuli. For example, the threat value of pain can be manipulated by telling participants performing a cold pressor task, “when feeling a tingling sensation in your hand, this may be the first signs of frostbite” [36] or warning patients that “lifting weights may cause back injury” [16], without actually experiencing these associations. But there is also a nonverbal pathway, during which the mere observation of another person in pain can be sufficient to install fear of that

particular stimulus [12]. In one study, participants observed human models who performed a cold pressor task, in which the color of the water (orange or pink) was associated with painful or neutral facial expressions [13]. When tested themselves, the observers’ fear and pain scores show that they learned the CS–pain associations that they previously observed in the model, despite equal temperatures of both cold pressors. An intriguing yet untested idea is that interactions among these pathways may facilitate learning. For example, previous observational learning may enhance subsequent experiential learning of pain-related fear during the actual encounter of a similar CS–pain pairing.

4. The role of (un)predictability

Although fear conditioning research and its application to pain-related fear has been valuable for the understanding of chronic regional musculoskeletal pain, it may not be appropriate for more generalized pain disorders. Presumably because of lack of apparent safety cues, experiencing an unpredictable threat induces a more general form of distress, demonstrated by feelings of worry and chronic apprehension [27]. The degree of predictability may have an impact on the experienced pain intensity. Indeed, an interesting

finding in the study using joystick movements was that in the unpredictable condition, both pain intensity and unpleasantness of the pain were rated significantly higher as compared with the paired condition, despite the equal number and physical intensities of the stimuli. The role of predictable versus unpredictable pain and the effects of different types of (un)predictability (pain duration, pain offset, pain location, pain quality, etc.) clearly warrants more systematic experimental scrutiny, and may open a novel window on the understanding of generalized pain syndromes.

5. Weighing pain and nonpain goals

Another concern with the current FA model relates to the idea that pain-related fear emerges in a context of multiple goals [6,37,38]. The goal to avoid pain is only one to be pursued in an environment with concomitant, often competing goals. Indeed, one of the most debilitating consequences of pain-related avoidance behavior is the withdrawal from previously valued activities. In this respect, chronic pain patients frequently weigh the value of their pain avoidance against the costs related to the loss of valued activities [31,32]. For example, in one study goal, self-efficacy, goal conflict, and pain severity independently predicted pain-induced fear, which in turn mediated the effects of goal conflict on physical disability and depression in chronic low back pain patients [18]. The idea that pain-related goal conflicts may increase the threat value of the pain is an intriguing one, largely left untested [19].

6. Novel assessment tools for pain-related fear

If pain-related fear mechanisms are a distinct trajectory by which acute pain becomes chronic, then they should be potent factors for identification. The Örebro Musculoskeletal Pain Screening Questionnaire was developed for this purpose, with which it is possible, with reasonable accuracy, to identify patients who risk developing persistent disability [14,25]. However, more research on the relative contribution of the FA variables on future outcome is needed to construct instruments with increased predictive accuracy. Also, the question remains regarding for whom, when, and where screening is best conducted, as well as how the information should be used in planning further assessment and treatment.

The challenge of assessment is to identify patients in whom pain-related fear and its ensuing avoidance are a significant problem. In our 2000 review, we called for broadening assessment techniques extending the cognitive aspects to the behavioral and physiological features. Today, advances have been made in self-report measures, but little progress in other aspects. Questionnaires have been further developed [9,30], alternative pictorial assessment methods are now available for identifying perceived “harmful” movements [22,34], and automated activity monitoring devices has been used [35]. One particular challenge is to objectively measure escape and avoidance behavior. Not just for the avoidance of fear-eliciting activities, but also for the more subtle safety-seeking behaviors, reliable and valid assessment methods are currently lacking [29].

7. The reduction of pain-related fear

Treatment procedures and the evaluation of their effects have developed dramatically. Exposure *in vivo* has a strong pedigree as one of the most powerful cognitive behavioral treatments for reducing disabling fear and anxiety, and has now been applied in patients with chronic pain [2,15]. Several single-subject experimental studies show impressive improvements in fear, catastrophizing, and function, which have initiated subsequent randomized controlled trials, (e.g., [23]). Although the results vary, effect sizes are at most moderate, suggesting room for

improvement. Recent studies on the mechanisms behind exposure therapy reveal that CS–US associations are not “unlearned” during the extinction of fear, but that instead, inhibitory responses are learned during extinction. This means that the original (excitatory) CS–US association remains intact, but competes with a new (inhibitory) CS–“no US” association. Exposure can thus best be designed such that new nonthreat associations be formed, and subsequently generalized across time and contexts. Such an approach shifts the focus toward the inclusion of multiple CSs, eliminating safety behaviors during exposure and using mental rehearsal to bridge exposure contexts to other daily life contexts [5]. Given these new insights, remaining issues call for further scrutiny. What is the relative impact of actual exposure versus observational or verbally transmitted information about associations between the CSs and the absence of pain or its feared consequences [24]? Do the effects of exposure to proprioceptive conditioned stimuli generalize to interoceptive ones, and vice versa [7]? What is the moderating effect of patient levels of executive control? How can fear-reduction techniques be applied in secondary prevention [28]? What are minimal competencies for treatment providers executing exposure? Many of these ways to optimize the effects of exposure await experimental and clinical testing.

8. Conclusion

The last decade has seen a surge in the study of the FA model of pain in both basic and clinical investigations. The recent literature mainly supports the basic assumptions of the model, but it also provides greater depth, inspiring future research and novel clinical applications. In particular, the model draws on the associative learning, and experimental research provides a fertile ground for future work on the intricacies of its mechanisms. Research is needed, for example, to clarify how interoceptive stimuli might work as a CS, what the role of (un)predictability has on fear responding, and how competing goals may influence fear learning. Clinically, the FA model has made a contribution to advance our understanding of the development and reduction of persistent disability, but considerable challenges remain in order to harvest the full benefit of the knowledge gained. Future efforts should focus on developing more specific assessment procedures that could direct clinicians to the best treatment options and optimize tailoring. Although exposure techniques are clearly helpful, there is promise in developing them further in order to understand how the model operates in patients as well as for more effective applications. These new avenues are likely to strengthen the predictive validity and clinical utility of future FA models in the context of chronic pain and disability.

Conflict of interest statement

The authors have no conflict of interest to report.

Acknowledgements

Johan W.S. Vlaeyen was supported by the Social Sciences Research Council of the Netherlands (NWO), and the Research Foundation, Flanders, Belgium (FWO Vlaanderen). Steven J. Linton received grants from the Swedish Council for Working Life and Social Research.

References

- [1] Arntz A, Claessens L. The meaning of pain influences its experienced intensity. *Pain* 2004;109:20–5.
- [2] Bailey KM, Carleton RN, Vlaeyen JWS, Asmundson GJG. Treatments addressing pain-related fear and anxiety in patients with chronic musculoskeletal pain: a preliminary review. *Cogn Behav Ther* 2010;39:46–63.
- [3] Beecher HK. Pain in men wounded in battle. *Ann Surg* 1946;123:96–105.

- [4] Bradley MM, Silakowski T, Lang PJ. Fear of pain and defensive activation. *Pain* 2008;137:156–63.
- [5] Craske MG, Kircanski K, Zelikowsky M, Mystkowski J, Chowdhury N, Baker A. Optimizing inhibitory learning during exposure therapy. *Behav Res Ther* 2008;46:5–27.
- [6] Crombez G, Eccleston C, Van Damme S, Vlaeyen JW, Karoly P. The fear avoidance model of chronic pain: the next generation. *Clin J Pain*; in press
- [7] De Peuter S, Van Diest I, Vansteenwegen D, Van den Bergh O, Vlaeyen JW. Understanding fear of pain in chronic pain: interoceptive fear conditioning as a novel approach. *Eur J Pain* 2011;15:889–94.
- [8] Dworkin BR, Dworkin S. Learning of physiological responses: II. Classical conditioning of the baroreflex. *Behav Neurosci* 1995;109:1119–36.
- [9] George SZ, Valencia C, Zeppieri G, Robinson ME. Development of a self-report measure of fearful activities for patients with low back pain: the Fear of Daily Activities Questionnaire. *Phys Ther* 2009;89:969.
- [10] Gheldof EL, Crombez G, Van den Bussche E, Vinck J, Van Nieuwenhuysse A, Moens G, Mairiaux P, Vlaeyen JW. Pain-related fear predicts disability, but not pain severity: a path analytic approach of the fear-avoidance model. *Eur J Pain* 2010;14:870. e871–870. e879.
- [11] Goubert L, Crombez G, Van Damme S. The role of neuroticism, pain catastrophizing and pain-related fear in vigilance to pain: a structural equations approach. *Pain* 2004;107:234–41.
- [12] Goubert L, Vlaeyen JW, Crombez G, Craig KD. Learning about pain from others: an observational learning account. *J Pain* 2011;12:167–74.
- [13] Helsen K, Goubert L, Peters ML, Vlaeyen JW. Observational learning and pain-related fear. An experimental study with colored cold pressor tasks. *J Pain* 2011;12:1230–9.
- [14] Hockings RL, McAuley JH, Maher CG. A systematic review of the predictive ability of the Örebro Musculoskeletal Pain Questionnaire. *Spine* 2008;33:E494–500.
- [15] Hollander M, de Jong JR, Volders S, Goossens MEJB, Smeets RJEM, Vlaeyen JWS. Fear reduction in patients with chronic pain: a learning theory perspective. *Exp Rev Neurother* 2010;10:1733–45.
- [16] Houben RM, Ostelo RW, Vlaeyen JW, Wolters PM, Peters M, Stomp-van den Berg SG. Health care providers' orientations towards common low back pain predict perceived harmfulness of physical activities and recommendations regarding return to normal activity. *Eur J Pain* 2005;9:173–83.
- [17] Jensen JN, Karpatschof B, Labriola M, Albertsen K. Do fear-avoidance beliefs play a role on the association between low back pain and sickness absence? A prospective cohort study among female health care workers. *J Occup Environ Med* 2010;52:85–90.
- [18] Karoly P, Okun MA, Ruehlman LS, Pugliese JA. The impact of goal cognition and pain severity on disability and depression in adults with chronic pain: an examination of direct effects and mediated effects via pain-induced fear. *Cogn Ther Res* 2008;32:418–33.
- [19] Karsdorp PA, Vlaeyen JW. Goals matter: both achievement and pain-avoidance goals are associated with pain severity and disability in patients with low back and upper extremity pain. *Pain* 2011;152:1382–90.
- [20] Konorski J. Integrative activity of the brain. An interdisciplinary approach. Chicago, IL: University of Chicago Press; 1967.
- [21] Leeuw M, Goossens ME, Linton SJ, Crombez G, Boersma K, Vlaeyen JW. The fear-avoidance model of musculoskeletal pain: current state of scientific evidence. *J Behav Med* 2007;30:77–94.
- [22] Leeuw M, Goossens ME, van Breukelen GJ, Boersma K, Vlaeyen JW. Measuring perceived harmfulness of physical activities in patients with chronic low back pain: the photograph series of daily activities-short electronic version. *J Pain* 2007;8:840–9.
- [23] Leeuw M, Goossens ME, van Breukelen GJ, de Jong JR, Heuts PH, Smeets RJ, Koke AJ, Vlaeyen JW. Exposure in vivo versus operant graded activity in chronic low back pain patients: results of a randomized controlled trial. *Pain* 2008;138:192–207.
- [24] Linton SJ, McCracken LM, Vlaeyen JW. Reassurance: help or hinder in the treatment of pain. *Pain* 2008;134:5–8.
- [25] Melloh M, Elfering A, Egli-Presland C, Roeder C, Barz T, Rolli-Salathé C, Tamcan O, Mueller U, Theis J. Identification of prognostic factors for chronicity in patients with low back pain: a review of screening instruments. *Int Orthop* 2009;33:301–13.
- [26] Meulders A, Vansteenwegen D, Vlaeyen JWS. The acquisition of fear of movement-related pain and associative learning: a novel pain-relevant human fear conditioning paradigm. *Pain* 2011;152:2460–9.
- [27] Mineka S, Kihlstrom JF. Unpredictable and uncontrollable events: a new perspective on experimental neurosis. *J Abnorm Psychol* 1978;87:256–71.
- [28] Nicholas MK, George SZ. Psychologically informed interventions for low back pain: an update for physical therapists. *Phys Ther* 2011;91:765–76.
- [29] Prkachin KM, Schultz IZ, Hughes E. Pain behavior and the development of pain-related disability: the importance of guarding. *Clin J Pain* 2007;23:270–7.
- [30] Roelofs J, van Breukelen G, Sluiter J, Frings-Dresen MHW, Goossens M, Thibault P, Boersma K, Vlaeyen JWS. Norming of the Tampa Scale for Kinesiophobia across pain diagnoses and various countries. *Pain* 2011;152:1090–5.
- [31] Roy M. Weighting pain avoidance and reward seeking: a neuroeconomical approach to pain. *J Neurosci* 2010;30:4185–6.
- [32] Schrooten MG, Vlaeyen JW. Becoming active again? Further thoughts on goal pursuit in chronic pain. *Pain* 2010;149:422–3.
- [33] Swinkels-Meewisse IE, Roelofs J, Oostendorp RA, Verbeek AL, Vlaeyen JW. Acute low back pain: pain-related fear and pain catastrophizing influence physical performance and perceived disability. *Pain* 2006;120:36–43.
- [34] Turk DC, Robinson JP, Sherman JJ, Burwinkle T, Swanson K. Assessing fear in patients with cervical pain: development and validation of the Pictorial Fear of Activity Scale-Cervical (PFAcS-C). *Pain* 2008;139:55–62.
- [35] Verbunt JA, Huijnen IP, Koke A. Assessment of physical activity in daily life in patients with musculoskeletal pain. *Eur J Pain* 2009;13:231–42.
- [36] Vlaeyen JW, Hanssen M, Goubert L, Vervoort T, Peters M, van Breukelen G, Sullivan MJ, Morley S. Threat of pain influences social context effects on verbal pain report and facial expression. *Behav Res Ther* 2009;47:774–82.
- [37] Vlaeyen JW, Morley S. Active despite pain: the putative role of stop-rules and current mood. *Pain* 2004;110:512–6.
- [38] Vlaeyen JWS, Crombez G, Linton SJ. The fear-avoidance model of pain: we are not there yet. Comment on Wideman et al. "A prospective sequential analysis of the fear-avoidance model of pain" [Pain, 2009] and Nicholas "First things first: reduction in catastrophizing before fear of movement" [Pain, 2009]. *Pain* 2009;146:222.
- [39] Vlaeyen JWS, Linton SJ. Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain* 2000;85:317–32.
- [40] Wideman TH, Adams H, Sullivan MJ. A prospective sequential analysis of the fear-avoidance model of pain. *Pain* 2009;145:45–51.